

# **Stable Boundary Layers and Pollution Transport in Mountain Terrain**

**Craig B. Clements**  
**Atmospheric Science Programme**  
**University of British Columbia**  
**Vancouver, Canada**

End-of-Summer Workshop  
GCEP  
2002

# **Overview**

**1) Introduction: Boundary Layer and Stability Basics**

**2) Why is mountain meteorology important?**

**3) Observations**

- **Yosemite National Park, Sierra Nevada**
- **Peter Sinks, Utah**
- **Monson Vineyard, WA**


**4) Summary**


# Air Pollution Meteorology- Basics

---

## Pollution Dispersion:

- Wind speed and Direction
- Plume rise
- Turbulence
- Atmospheric Stability





# What is Mountain Meteorology?

---



**Tioga Pass, CA  
Winter 1995**

# Lee Vining Canyon, eastern Sierra Nevada

---





## Pilot Balloon Release- LVC, Sierra Nevada

---



# No-Lift Balloons at Tioga Pass, Sierra Nevada

---






# Why is mountain meteorology important?



**The Western US**

# Mountain Meteorology: Applications for Air Pollution

Daytime- Late Afternoon



---

## Early Evening

Synoptic-Scale Winds



Valley Winds




Drainage  
Flows



Cold Air Pool









# Tethersonde in Yosemite Valley, Sierra Nevada

---



# Vertical Profiles – Yosemite Valley (March 16, 1998)



# Campfire smoke plume – Cayuse Creek, ID

---




# Polluted CBL: Coast Mountains, British Columbia

---





# Concepts of cold pool buildup



# Valley/basin heat budget


$$\underbrace{\bar{\rho} c_p \frac{f\theta}{ft} dv}_A = \underbrace{-\bar{\rho} c_p \overline{(\mathbf{V}\theta)} dv}_B + \underbrace{-\frac{\theta}{T} \overline{\mathbf{R}} dv}_C + \underbrace{-\bar{\rho} c_p \overline{(\mathbf{V}\theta)} dv}_D$$

**Term A** : Rate of change of potential temperature (**Heat Storage**)


**Term B**: convergence of potential temperature flux by mean wind (**Advection**)

**Term C**: convergence of **Radiative Flux**

**Term D**: convergence of **Turbulent Sensible Heat Flux**



# Mass conservation - valleys vs basins




# The Peter Sink - limestone sinkhole

---







# Map of Peter Sinks



# Instrument locations



# Peter Sinks ridgeline



# Operations Base Tent

---






# Tethersondes

---



# Tethersonde, 3D sonic and net radiometer






# CSI weather station & thermocouple mast

---




# Temperature time series






# Net radiation and sensible heat flux




# Potential temperature profiles at TS1




# Accumulated Heat Losses from Peter Sinks

$$S = - \frac{\int_0^h c_p \rho(z) [\theta_0 - \theta(z)] A(z) dz}{A_h}$$




# Synchronous temperature profiles





# Observed Slope Flow Structure – Peter Sinks



## Summary- from Peter Sinks Experiments

---

- Air over basin sidewall became warmer (1-5 K) than air over basin center- does not support continued slope flows.
- Formation of persistent superadiabatic layers within one of the most stable atmospheric structures on Earth!
- Downslope flows played minor role in cold pool formation.
- Heat loss from basin atmosphere is initially at the net rate of longwave loss, but decreases with time through night.

# Monson Ranch Vineyard, WA

---

